THE GENERAL ASSEMBLY OF PENNSYLVANIA

HOUSE RESOLUTION

No. 719

Session of 2010

INTRODUCED BY FREEMAN, HENNESSEY, ADOLPH, BAKER, BEYER, BISHOP, BOBACK, BRADFORD, BRENNAN, BRIGGS, BROOKS, BUXTON, CALTAGIRONE, CAUSER, CLYMER, COHEN, CONKLIN, D. COSTA, CUTLER, DALEY, DENLINGER, DIGIROLAMO, EVERETT, FAIRCHILD, GEIST, GEORGE, GIBBONS, GILLESPIE, GODSHALL, GOODMAN, GRELL, GRUCELA, HARHART, HELM, HESS, HICKERNELL, KIRKLAND, KORTZ, LONGIETTI, MAHER, MAHONEY, MAJOR, MANN, MARSICO, McGEEHAN, MCILVAINE SMITH, MILLARD, MURPHY, MUSTIO, MYERS, M. O'BRIEN, PALLONE, PASHINSKI, PAYTON, PHILLIPS, PICKETT, READSHAW, REICHLEY, ROCK, ROEBUCK, ROSS, SAINATO, SAMUELSON, SANTONI, SAYLOR, SCAVELLO, SEIP, SIPTROTH, SOLOBAY, SONNEY, STABACK, STERN, TALLMAN, TRUE, VULAKOVICH, WATSON, HAHN, SWANGER AND PAYNE, SEPTEMBER 7, 2010

INTRODUCED AS NONCONTROVERSIAL RESOLUTION UNDER RULE 35, SEPTEMBER 7, 2010

A RESOLUTION

- Designating the month of September 2010 as "Colorectal Cancer Examination Awareness Month" in Pennsylvania.
- 3 WHEREAS, Colorectal cancer is both the third most common
- 4 cancer diagnosis and the second leading cause of cancer death in
- 5 the United States of America; and
- 6 WHEREAS, A family history of colorectal cancer or a personal
- 7 history of an inflammatory bowel disease is considered to be
- 8 among the biggest risk factors for the disease, with other risk
- 9 factors including obesity, alcohol consumption, red and
- 10 processed meat consumption and diabetes; and
- 11 WHEREAS, Men are 35% more likely to get colorectal cancer
- 12 than women and African Americans are 20% more likely to get this

- 1 cancer than other races or ethnicities; and
- 2 WHEREAS, It is estimated that as many as 60% of colorectal
- 3 cancer deaths could be prevented if all men and women 50 years
- 4 of age or older were screened routinely, for colorectal
- 5 examinations save lives; this is because colorectal cancer
- 6 develops from precancerous polyps in the colon and rectum which
- 7 can be seen in examinations and removed before they turn into
- 8 cancer; and
- 9 WHEREAS, Screening tests also can find colorectal cancer
- 10 early, when treatment works best; and
- 11 WHEREAS, Since the 1990s, colorectal cancer incidence and
- 12 mortality rates have been declining and the five-year survival
- 13 rate has been increasing; and
- 14 WHEREAS, Early colon cancer detection through sigmoidoscopies
- 15 and, where indicated, colonoscopies has contributed to the
- 16 decline in the death rate; and
- 17 WHEREAS, Use of these examinations and good dietary habits
- 18 should be encouraged through awareness programs; therefore be it
- 19 RESOLVED, That the House of Representatives designate the
- 20 month of September 2010 as "Colorectal Cancer Examination
- 21 Awareness Month" in Pennsylvania.